

Elliptic flow and the high p_T ridge in Au+Au collisions

N. K. Pruthi (for the STAR collaboration)
Panjab University, Chandigarh-160014, India*

In this paper we look for a coupling between intermediate p_T particle pairs and the v_2 of the remaining low p_T particles. We find that the shape of the flow vector distribution, which is calculated from all low p_T tracks, depends in a non-trivial way on the angular separation between the high p_T particles in the event. Our analysis is based on 200 GeV Au+Au collisions measured with the STAR

1. Introduction

Dihadron correlations are used in heavy-ion collisions to study the effect of jet-quenching and other interactions of jets with the medium created in the collision overlap zone [1, 2]. In non-central collisions, this overlap zone is not symmetric. The spatial asymmetry leads to large azimuthal anisotropies in momentum space resulting in high values of the second coefficient in the Fourier decomposition of the azimuthal distribution of produced particles with respect to reaction plane, (v_2) [3]. Dihadron measurements from STAR have revealed a correlation structure which is narrow in the azimuthal direction and broad in longitudinal direction (the narrow $\Delta\phi$, broad $\Delta\eta$ “ridge”) [4, 5, 6]. This structure is unique to nucleus-nucleus collisions and its amplitude shows a non-monotonic rise with increase in collision centrality [7]. We search for evidence of jet-interactions with the medium by studying v_2 for events containing correlated dihadron pairs.

2. Method and discussion

Our analysis is based on 16 million Au+Au at 200GeV center of mass energy measured with the STAR detector [8]. We select events with at least two tracks having $p_T > 2.0$ GeV and calculate the $|q|$ for those events. The q (“flow”) vector can indicate if particles are produced in preferential direction. $|q|$ is calculated from $q_x = \Sigma \cos(2\phi_i)$ and $q_y = \Sigma \sin(2\phi_i)$ [9], where the sums run over all tracks having $p_T < 2.0$ GeV/c and ϕ_i is azimuthal angle of track w.r.t reaction plane. The distribution of $|q|$ ($dN/d|q|$) depends on the number of particles, v_2 , and other correlations not related to v_2 [10]. If there are more than two tracks with $p_T > 2.0$ GeV/c, we select the leading and sub-leading tracks. This way we select one dihadron pair per event. This allows us to associate the $|q|$ of an event with the angular separation of a high momentum pair. In this way we can look for modifications to $dN/d|q|$ for events with or without correlated high p_T pairs.

Fig. 1 (left) shows the distribution of the relative angle between leading and sub-leading dihadron pairs with $p_T > 2.0$ GeV/c. If v_2 were the only source of correlations then $dN/d\Delta\phi$ should have the following shape: $C(\Delta\phi) = b_0(1 + 2\langle v_2^{trig} v_2^{asso} \rangle \cos(2\Delta\phi))$, where v_2^{trig} and v_2^{asso} are the v_2 values respectively, for the leading and subleading (trigger and associate) particles. Other physical processes leading to correlations are particle decays, fragmentation of partons (jets), and HBT. Jet correlations are expected to be prominent for high p_T dihadron pairs so a two-component model is used to fit $dN/d\Delta\phi$: a “jet” component on top of a v_2 modulated “background” [1, 2, 11]. Assuming the two components model is applicable, we can extract the fraction of correlated and uncorrelated pairs for each $\Delta\phi$ bin.

*For the full list of STAR authors and acknowledgements, see appendix ‘Collaborations’ of this volume

The next step of our analysis is to study $dN/d|q|$ for the events with high p_T pairs as a function of the angle $\Delta\phi$ and $\Delta\eta$ of the high p_T pairs. If jets are the source of the correlations (Signal) in Fig. 1 and they interact with the medium, they may modify the v_2 of the event. That modification would then change the shape of $dN/d|q|$. Or alternatively, if the correlations beyond the v_2 modulation in Fig. 1 is caused by the same space-momentum correlation that give rise to v_2 , then the correlation signal may result to large values of v_2 [12]. This would also cause $dN/d|q|$ to depend on $\Delta\phi$ and/or $\Delta\eta$. Fig. 1 (right) shows average magnitude of flow vector ($\langle|q|\rangle$) vs. $\Delta\phi$ of the leading and subleading hadrons for either $|\Delta\eta| > 0.7$ or $|\Delta\eta| < 0.7$. We observe a non-trivial dependence on $\Delta\phi$. No prominent $\Delta\eta$ dependence is seen in this figure.

We also studied $\langle p_T \rangle$ vs. $\Delta\phi$ and find that within errors it is independent of the angle between the high p_T tracks. The multiplicity distribution for the events with two tracks with $p_T > 2.0$ GeV only deviates from the inclusive sample for peripheral events.




Figure 1: (left) $dN/d\Delta\phi$ for triggered events and the background function. (right) Non-trivial behavior of $\langle q \rangle$ with $\Delta\phi$ for ridge and ridge+jet events

Having observed a non-trivial dependence of $\langle q \rangle$ on $\Delta\phi$, we next attempt to convert this into a v_2 for the events that had a correlated pair. We categorize the events in two parts depending on the highest p_T particles (1) Ridge events : $|\Delta\eta| > 0.7$, (2) Ridge+jet events : $|\Delta\eta| < 0.7$ and studied the $dN/d|q|$ distributions.

We calculate the q-distributions for the signal and background by dividing the near-side into two bins ($\Delta\phi < 4\pi/20$ and $4\pi/20 < \Delta\phi < 8\pi/20$) and solving equations: $(S_1 + B_1) * dN_1 = S_1 * dN_S + B_1 * dN_B$, $(S_2 + B_2) * dN_1 = S_2 * dN_S + B_2 * dN_B$. $S_{1,2}$ is the number of events yielding a correlated pair in bin 1 or 2. $B_{1,2}$ is the number of events yielding an un-correlated pair in bin 1 or 2. dN_1 and dN_2 are the q-distributions for events in bin 1 and 2. dN_S and dN_B are the q-distributions for events giving correlated pairs (signal) and uncorrelated pairs (background). dN_1 and dN_2 are measured. $S_{1,2}$ and $B_{1,2}$ are extracted from the $dN/d\Delta\phi$ distributions by applying the two-component model. Then dN_S and dN_B can be determined from the two equations above. Elliptic flow is calculated by fitting the q-distributions (Fig. 2(left)) for signal for ridge and ridge+jet events. Two parameters can be extracted from the q-distribution : $(\langle v_2 \rangle^2 - \sigma_{v_2}^2) = v_2 \{q\}^2$ and $(\delta_2 + 2\sigma_{v_2}^2) = \sigma_{dyn}^2$. $\delta_2 = \langle \cos(2(\phi_1 - \phi_2)) \rangle - v_2^2$. The term accounts for correlations that are not related to the reaction plane and σ_{v_2} is the rms width of the v_2 distribution.

In Fig. 2(right), we report $v_2\{2\}$ for the event classes defined above. We use the ansatz that the near-side correlation contains a jet part (narrow in $\Delta\phi$ and $\Delta\eta$) and a ridge part (narrow in $\Delta\phi$ but broad in $\Delta\eta$) [6]. By calculating the signal v_2 for ridge and ridge+jet events we calculated v_2 for jet events by using equations

$$Area^{ridge+jet} v_2^{ridge+jet} = Area^{ridge} v_2^{ridge} + Area^{jet} v_2^{jet} \quad (1)$$

$$Area^{jet} = Area^{ridge+jet} - Area^{ridge} * Acceptance\ factor \quad (2)$$

$$Acceptance\ factor = \frac{Area^{ridge+jet(background)}(\Delta\eta < 0.7)}{Area^{ridge(background)}(\Delta\eta > 0.7)} \quad (3)$$




Figure 2: (left) q -vector distribution fit for events yielding correlated pair(signal)(right) Comparison of v_2 for jet signal with ridge signals

This allows us to project the ridge from $|\Delta\eta| > 0.7$ to $|\Delta\eta| < 0.7$. The jet is the correlation that remains.

The selection of one unique pair per event, the application the two-component model, and the simple algebra above allows us to calculate $v_2\{q\}^2 = v_2^2 - \sigma_{v_2}^2$ and $\sigma_{q,dyn}^2 = \delta_2 + 2\sigma_{v_2}^2$ for events giving rise to pairs of particles in the jet-cone region, in the ridge region, and in the background. The values of $v_2\{q\}$ vs. collision impact parameter b are presented in Fig. 2(right). $v_2\{q\}$ for the events that had a high p_T pair contributing to the ridge-like correlation exhibit a slightly larger $v_2\{q\}$ for non-central and peripheral collisions than the corresponding events contributing a pair to the background. We also attempt to determine if the $v_2\{q\}$ values for events contributing pairs to the jet-like correlation are larger or smaller than those contributing to the background. Within the current estimates of the systematics uncertainties the values are consistent with v_2 from inclusive events.

3. Summary

We have developed an analysis to search for jet interactions with the medium and/or possible effects linking the high p_T ridge correlation to v_2 . The latter could occur if the same physics mechanism underlies both phenomena as proposed in several references [13, 14, 15, 16]. By selecting one high- p_T pair per event and the two component model, we find that $v_2\{q\}$ calculated from $dN/d|q|$ for events yielding pairs in the ridge is slightly larger v_2 for non-central and peripheral collisions than the events yielding pairs in the background.

References

- [1] C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **90**, 082302 (2003).
- [2] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **95**, 152301 (2005).
- [3] K. H. Ackermann *et al.* [STAR Collaboration], Phys. Rev. Lett. **86**, 402 (2001); J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 052302 (2004); C. Adler *et al.* [STAR Collaboration], Phys. Rev. C **66**, 034904 (2002); J. Adams *et al.* [STAR Collaboration], Phys. Rev. C **72**, 014904 (2005).
- [4] J. Adams *et al.* [STAR Collaboration], Phys. Rev. C **73**, 064907 (2006); J. Adams *et al.* [STAR Collaboration], J. Phys. G **32**, L37 (2006).
- [5] J. Adams *et al.* [Star Collaboration], Phys. Rev. C **75**, 034901 (2007).
- [6] J. Putschke, J. Phys. G **34**, S679 (2007).
- [7] M. Daugherty, QM08 Proceedings;
- [8] K. H. Ackermann *et al.* [STAR Collaboration], Nucl. Instrum. Meth. A **499**, 624 (2003).
- [9] J. Y. Ollitrault, arXiv:nucl-ex/9711003; A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C **58**, 1671 (1998); N. Borghini, P. M. Dinh and J. Y. Ollitrault, Phys. Rev. C **63**, 054906 (2001).

- [10] P. Sorensen [STAR Collaboration], arXiv:0808.0356 [nucl-ex].
- [11] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **97**, 052301 (2006).
- [12] S. A. Voloshin, Phys. Lett. B **632**, 490 (2006).
- [13] S. Mrowczynski, Phys. Lett. B **214**, 587 (1988) [Erratum-ibid. B **656**, 273 (2007)].
- [14] A. Majumder, B. Muller and X. N. Wang, Phys. Rev. Lett. **99**, 192301 (2007).
- [15] R. C. Hwa and C. B. Yang, arXiv:0801.2183 [nucl-th].
- [16] A. Dumitru, F. Gelis, L. McLerran and R. Venugopalan, arXiv:0804.3858 [hep-ph].